

A Report of Integrated Vector Surveillance in Dhangadhi Sub-Metropolitan City-12, Kailali

(February 2024)



Submitted by:

Hem Raj Joshi

VCSO, PHD Doti

Prakriti Pant

Entomologist, PHD Doti

Submitted to:

Sudurpaschim Province

Ministry of Social Development

Provincial Health Directorate

Rajpur, Dipayal, Doti

2024

ACKNOWLEDGEMENT

We would like to express our sincere gratitude to Health Directorate, Rajpur, Doti for providing us such an opportunity to conduct “Integrated Vector Surveillance in Dhangadhi Sub-Metropolitan City, Kailali”. We are thankful to our team members and local citizens for their dedication and assistance during this work.

Team members:

1. Hem Raj Joshi (Vector Control Supervisor Officer, PHD, Doti)
2. Dipendra Singh Rawal (Vector Control Supervisor Officer, Mahakali Provincial Hospital, Kanchanpur)
3. Prakriti Pant (Entomologist, PHD, Doti)
4. Bhuwaneshwari Joshi (Insect Collector, PHD, Doti)
5. Rachana Bhatta Paneru (Insect Collector, PHD, Doti)

Hem Raj Joshi

Prakriti Pant

TABLE OF CONTENTS

ACKNOWLEDGEMENT	ii
ABSTRACT	iv
1. INTRODUCTION	1
2. MATERIALS AND METHODS	4
2.1. Study Area	4
2.2. Materials	5
2.3. Vector Sampling and Identification	5
2.4. Data Analysis	6
3. RESULTS	7
3.1. Species composition of mosquito vectors	7
3.2. Indoor resting collection	7
3.3. Light trap collection	8
3.4. Larval collection	9
4. DISCUSSIONS	10
5. CONCLUSIONS AND RECOMMENDATIONS	10
REFERENCES	11
PHOTOPLATES	14
Annex: 1	16
Annex: 2	16
Annex: 3	17

ABSTRACT

Vector borne diseases are one of major public health problems in Sudurpashchim Province of Nepal. Entomological surveillance enables us to understand composition, abundance and distribution of vector species in any locality, and is an essential component of malaria vector control programs, operational activities and research. Dhangadhi Sub-Metropolitan City of Kailali district lies in the southern part of the district and connected to India border as well. The present survey was conducted during the first week of February 2024 to know the current status of arthropod vectors in this area. A total of 111 specimens of mosquito belonging to seven species were collected and identified from the study area. *Culex quinquefasciatus* was found to be most abundant (73.87%) followed by *Anopheles fluviatilis* (12.61%), *Anopheles annularis* (6.31%) and others. *Anopheles culicifacies*, *Culex tritaeniorhynchus*, *Culex jacksoni* and *Armigeres subalbatus* were also recorded in this study. There were found vector species for malaria, filariasis and Japanese Leishmaniasis, so it is important to focus on management of breeding habitats and minimizing human-vector contact in this area.

Key Words: Kailali, Dhangadhi, Anopheles, Culex, Armigeres

1. INTRODUCTION

Insects belong to the largest phylum in the animal kingdom, arthropoda, and play key roles in ecosystem. Yet, blood-feeding i.e. hematophagous arthropods such as mosquitoes, sandflies, ticks, mites etc. serve as vectors for devastating parasitic and viral human diseases and show tremendous individual variation in their capacity to transmit diseases (Baxter et al., 2017). Vector-borne diseases (VBDs) are caused by parasites, viruses and bacteria and transmitted to humans by different vectors. VBDs are major public health problems particularly in tropical and sub-tropical regions of the world. In the WHO South-East Asia Region (SEAR), the VBDs that are of major public health importance are malaria, dengue, Japanese encephalitis (JE), chikungunya, lymphatic filariasis and kala-azar (WHO, 2014). In addition to these, scrub typhus has been a major public health problem in Nepal since 2015 (Dhimal et al., 2021).

Mosquitoes are medically important insects as they act as vectors for major prevalent vector borne diseases in Nepal such as Malaria, Dengue, Lymphatic filariasis and Japanese encephalitis. Mosquitoes are being searched and studied in Nepal from 1958 in connection with "Malaria Eradication Program". Malaria was such a disaster back then in Nepal, so to reduce its burden different program were launched by the Government. First Malaria Control project was launched in 1954 with the support from USAID then National Malaria Eradication Program in 1958 again Malaria Control Program in 1978. To address perennial occurrence of malaria in foot hills, inner terai, valleys, and hard core forests of both terai and hills Roll Back Malaria (RBM) Program was launched in 1998 following the call of WHO to revamp the program. Nepal is working currently in Malaria Elimination Program 2014-2025 with the goal of malaria free Nepal by 2025 (EDCD, 2018). The high risk of acquiring the disease is attributed to the abundance of vector mosquitoes, mobile and vulnerable population, relative inaccessibility of the area, suitable temperature, environmental and socio-economic factors. Since presence and abundance of vector species affects disease burden proportionally, so National Malaria

Strategic Plan (2014-2025) endorsed vector surveillance as core intervention to reduce and interrupt the malaria transmission (EDCD, 2020a).

Malaria is a complex disease caused by protozoan parasites (Genus: *Plasmodium*) and transmitted by blood-feeding infectious *Anopheles* mosquitoes (Dutta & Dutt, 1978). Mosquitoes are bilaterally symmetrical belonging to the family Culicidae. These are delicate, slender flies; legs are two or three times as long as body, antennae filiform. Body is strongly humped; both head and abdomen drooping downwards. Larvae and pupae are aquatic. A total of 3,719 extant species of Culicidae are currently formally recognized worldwide (Mosquito Taxonomic Inventory, 2023). About 537 are species of *Anopheles* (Harbach, 2013) were reported and only 70-80 are known to transmit human malaria worldwide. Of these, 41 are considered to be the dominant vector species and capable of transmitting malaria by large (Sinka et al., 2012).

In 1990, Darsie and Pradhan published an extensive account of the mosquitoes of Nepal, recording 130 species in 14 genera. A total of 24 species of *Aedes* mosquitoes were reported from Gandaki, Janakpur, Mahakali, Seti, Koshi, Bagmati, Narayani and Karnali Zones of Nepal. This genus has been reported from low to high altitudes of Jumla district. The recorded 45 species of *Anopheles* mosquitoes are distributed from the Terai and inner Terai areas to the hills and mountains of the country. Among 45 *Anopheles* species only seven have been reported as malaria vectors of primary importance. These include: *Anopheles minimus*, *An. fluviatilis*, *An. annularis*, *An. maculatus*, *An. dravidicus*, *An. pseudowillmori*, and *An. willmori*. After *Anopheles* another largest genus of mosquito distributed in Nepal is *Culex* with 29 recorded species. Like *Anopheles*, *Culex* are also distributed throughout the country except high mountains but has been reported in the high altitudes of Jumla from various habitats. Over the last decades, deforestation and effective malaria control program using Dichloro- Diphenyl Trichloroethane (DDT) eliminated *An. minimus* during 1960s (Parajuli et al, 1981). *An. fluviatilis* is now the primary malaria vector in Nepal, *An. annularis* the secondary malaria vector and the *An. maculatus* complex members are seasonal malaria vectors in the mountain region of Nepal (Darsie & Pradhan, 1990; Rana, K.J., 2001).

Malaria has remained as a major public health problem impacting on the health and lives of a large proportion of people particularly in remote areas with low economies and concentrated along international borders (Dhimal et al., 2014a). As entomological surveillance can be defined as the regular, systematic collection, analysis and interpretation of entomological data for risk assessment, planning, implementation, monitoring and evaluation of vector control interventions; all the surveillance activities must be clearly linked to programme decisions to ensure optimal vector control (EDCD, 2020b). Effective vector control is reliant on knowledge of local vector species and their susceptibility to insecticides, as well as on vector and human behaviours that may allow mosquitoes to avoid contact with interventions and thereby maintain residual transmission (Afrane et al., 2005). It is well established that effective vector control programmes can make a major contribution towards advancing human and economic development. Vector control interventions have one of the highest returns on investment in public health. Aside from direct health benefits, reductions in vector-borne diseases will enable greater productivity and growth, reduce household poverty, increase equity and women's empowerment, and strengthen health systems (EDCD, 2020b).

Malaria is a significant public health concern in Nepal, especially during the monsoon season when mosquito breeding is more prevalent. The government and various organizations have been working to control malaria transmission through vector control measures, improved diagnosis, and timely treatment. Periodic collection of vector data is essential to inform vector control strategies and track their impact on malaria transmission. According to malaria micro-stratification- 2023, three provinces are under moderate to high risk. These provinces collectively have 10 high risk wards and 35 moderate risk wards (EDCD, 2024). The present integrated vector surveillance was carried out to investigate arthropod vector population of Dhangadhi sub-metropolitan city ward number 12. This study will be helpful in launching vector and disease control strategies in future.

2. MATERIALS AND METHODS

2.1. Study Area



Figure: 1. Map of Study Area

Kailali district of Nepal is located in Sudurpaschim Province having the spatial extent between latitude 28.390°N to 29.066°N and longitude 80.471°E to 81.294°E, covering an area of 3235 km². Dhangadi sub-metropolitan is surrounded by Kailari RM in east, Kanchanpur district separated by Mohana River in west, Godawai and Gauriganga Municipalities in North and India Border in South. It has 19 ward in total and most of the area is plain terai. Dhangadhi Sub-Metropolitan City-12 was chosen for vector surveillance.

2.2. Materials

- ✓ Aspirators
- ✓ Hand Torch
- ✓ Light Trap
- ✓ Plastic Cups
- ✓ Dipper and Dropper
- ✓ Vials
- ✓ Cotton, Mask, Gloves
- ✓ Chloroform
- ✓ Petri dishes
- ✓ Stereoscope

2.3. Vector Sampling and Identification

The entomological survey was performed on first week of February 2024. Adult mosquitoes/sandflies were collected from collection six human dwellings and four cattle shed, and the collecting sites were selected and fixed for everyday collection. For outdoor collection natural shelters such as tree holes, bushes near cattle shed were inspected. Indoor and outdoor resting mosquitoes were collected using torchlights and aspirators in the morning. Light trap was placed inside most suitable cattle shed among the sheds

selected for morning collection. GPS points for each sampling sites were recorded (Annex 3) along with temperature and humidity.

Oral informed consent was taken from each household before starting the collection of mosquitoes either in houses or in animal shelters. In cases where household member disagreed, the house was excluded from the collection plan and the immediate next one was chosen for study.

All the collected samples were brought to Entomology Laboratory of Province Health Directorate at Mahendranagar, Kanchanpur. Immature stages were kept for rearing in laboratory conditions. Adult mosquitoes, both collected and emerged from larval collection, were immobilized using Chloroform and identified using relevant taxonomic keys (Darsie & Pradhan, 1990; WHO, 2020)

2.4. Data Analysis

Data were managed in Microsoft Excel 2013, analysed and interpreted using its tools.

3. RESULTS

3.1. Species composition of mosquito vectors

A total of 111 specimens of mosquito belonging to seven species were collected and identified from the study area. Mosquitoes were collected through indoor resting collection, light trap collection and larval collection methods. Three species of *Anopheles*, three species of *Culex* and one species of *Armigeres* were recorded (Table 1). *Culex quinquefasciatus* was found to be most abundant (73.87%) followed by *Anopheles fluviatilis* (12.61%), *Anopheles annularis* (6.31%) and others. *Anopheles culicifacies*, *Culex tritaeniorhynchus*, *Culex jacksoni* and *Armigeres subalbatus* were also recorded in this study. Both male and female individuals were recorded and counted for analysis.

Table 1: Species composition of Vectors in Dhangadhi sub-metropolitan city

S.N.	Species	no. of Individuals	Relative Abundance (%)	Collection Method
1	<i>Anopheles fluviatilis</i>	14	12.61	AIRC, HIRC, LT, LC
2	<i>Anopheles annularis</i>	7	6.31	AIRC, HIRC, LT
3	<i>Anopheles culicifacies</i>	1	0.90	AIRC
4	<i>Culex quinquefasciatus</i>	82	73.87	AIRC, HIRC, LT
5	<i>Culex tritaeniorhynchus</i>	2	1.80	LT
6	<i>Culex jacksoni</i>	3	2.70	LC
7	<i>Armigeres subalbatus</i>	2	1.80	AIRC, LT
	Total	111		

3.2. Indoor resting collection

Total 77 mosquito specimens were collected through indoor resting collection from both animal and human dwellings. *Culex quinquefasciatus* was the most dominant mosquito species in animal as well as in human dwellings followed by *Anopheles fluviatilis*. In animal dwellings, *Culex quinquefasciatus* was most abundant species followed by *Anopheles fluviatilis*, *Anopheles annularis*, *Anopheles culicifacies* and *Armigeres subalbatus*. In human dwellings *Culex quinquefasciatus* was the most abundant followed by *Anopheles fluviatilis* and *Anopheles annularis* (Figure 2). Further abdominal condition

of indoor resting female mosquitoes was also investigated and most individuals were found to be fulfed (Annex 1).




Figure 2: Abundance of vector species through indoor resting collection

3.3. Light trap collection

In light trap 30 specimens were collected, where *Culex quinquefasciatus* was most abundant species followed by *Anopheles fluviatilis*, *Culex tritaeniorhynchus*, *Anopheles annularis* and *Armigeres subalbatus*. Many male specimens of *Culex quinquefasciatus* were also collected through light trap (Figure 3). Abdominal conditions of collected specimens were also investigated and most of them were found to be unfed (Annex 2).




Figure 3: Abundance of vector species through indoor resting collection

3.4. Larval collection

Four specimens were emerged from larval collections, of them 2 specimens were male *Culex jacksoni*, one was female *Culex jacksoni* and one specimen of female *Anopheles fluviatilis*.

4. DISCUSSIONS

Seven species of mosquitos were collected and identified during present study. Among the malaria vectors *Anopheles fluviatilis* and *Anopheles annularis* were recorded in this study. As from previous studies, *An. fluviatilis* is the primary vector species in Nepal, where as *An. annularis* is a secondary vector and *An. maculatus* complex is considered as seasonal vector (Dhimal et al., 2014b). Vectors of malaria has been reported from different geo-ecological areas of Nepal (Dhimal et al., 2014c; Pradhan et al., 1970) and in this study as well two malaria vectors were recorded. Further *Culex quinquefasciatus* and *Culex tritaeniorhynchus* were recorded from the study area, which are vectors for filariasis and Japanese Encephalitis respectively.

5. CONCLUSIONS AND RECOMMENDATIONS

The vectors for malaria, filariasis and Japanese Encephalitis were recorded during this study so the management of breeding sources and minimizing human-vector contact will be important measure to reduce the chances of disease outbreak at the study area.

REFERENCES

- Afrane, Y. A., Lawson, B. W., Githeko, A. K., & Yan, G. (2005). Effects of Microclimatic Changes Caused by Land Use and Land Cover on Duration of Gonotrophic Cycles of *Anopheles gambiae* (Diptera: Culicidae) in Western Kenya Highlands. *Journal of Medical Entomology*, 42(6), 974–980.
<https://doi.org/10.1093/jmedent/42.6.974>
- Baxter, R. H. G., Contet, A., & Krueger, K. (2017). Arthropod Innate Immune Systems and Vector-Borne Diseases. *Biochemistry*, 56(7), 907–918.
<https://doi.org/10.1021/acs.biochem.6b00870>
- Darsie, R. F., & Pradhan, S. P. (1990). The mosquitoes of Nepal: Their identification, distribution and biology. *Mosquito Systematics*, 22(2), 69–130.
- Dhimal, M., Ahrens, B., & Kuch, U. (2014a). Malaria control in Nepal 1963–2012: Challenges on the path towards elimination. *Malaria Journal*, 13(1), 241.
<https://doi.org/10.1186/1475-2875-13-241>
- Dhimal, M., Ahrens, B., & Kuch, U. (2014b). Species composition, seasonal occurrence, habitat preference and altitudinal distribution of malaria and other disease vectors in eastern Nepal. *Parasites & Vectors*, 7(1), 540. <https://doi.org/10.1186/s13071-014-0540-4>
- Dhimal, M., Ahrens, B., & Kuch, U. (2014c). Species composition, seasonal occurrence, habitat preference and altitudinal distribution of malaria and other disease vectors in eastern Nepal. *Parasites & Vectors*, 7, 540. <https://doi.org/10.1186/s13071-014-0540-4>

- Dhimal, M., Dumre, S. P., Sharma, G. N., Khanal, P., Ranabhat, K., Shah, L. P., Lal, B. K., Jha, R., Upadhyaya, B. P., Acharya, B., Shrestha, S. K., Davidson, S. A., Charoensinphon, P., & Karki, K. B. (2021). An outbreak investigation of scrub typhus in Nepal: Confirmation of local transmission. *BMC Infectious Diseases*, 21(1), 193. <https://doi.org/10.1186/s12879-021-05866-6>
- EDCD. (2020a). *National Malaria Strategic Plan (updated)*.
- EDCD. (2018). *Nepal Malaria Strategic Plan 2014-2025*.
<https://km.mohp.gov.np/document/%02%03%04%05%06%07%08%05%06%05nepal-malaria-strategic-plan-2014-2025>
- EDCD. (2020b). *National Guideline on Integrated Vector Management 2020*. EDCD.
<https://www.edcd.gov.np/resource-detail/national-guideline-on-integrated-vector-management-2020-new>
- EDCD. (2024). *Malaria Elimination Program*. EDCD.
<https://www.edcd.gov.np/section/malaria-elimination-program>
- Harbach, R. E. (2013). The Phylogeny and Classification of Anopheles. In *Anopheles mosquitoes—New insights into malaria vectors*. IntechOpen.
<https://doi.org/10.5772/54695>
- Hiran Dutta, M., & Ashok Dutt, K. (1978). Malarial ecology: A global perspective. *Social Science & Medicine. Part D: Medical Geography*, 12(2), 69–84.
[https://doi.org/10.1016/0160-8002\(78\)90010-2](https://doi.org/10.1016/0160-8002(78)90010-2)
- Mosquito Taxonomic Inventory. (2023). *Valid Species*. <https://mosquito-taxonomic-inventory.myspecies.info/valid-species-list>

- Pradhan, J. N., Shrestha, S. L., & Vaidya, R. G. (1970). Malaria Transmission In High Mountain Valleys Of West Nepal Including First Record Of Anopheles Maculatus Willmori (James) As A Third Vector Of Malaria. *Journal of Nepal Medical Association*, 8(3), Article 3. <https://doi.org/10.31729/jnma.1384>
- Rana, K.J. (2001). History of Malaria and Malaria control in Nepal. *Aravali Printers and Publishers P. Ltd. New Delhi*.
https://scholar.google.com/scholar_lookup?title=History%20of%20Malaria%20and%20Malaria%20Control%20in%20Nepal&publication_year=2001&author=Rana%2CKJ#d=gs_cit&t=1683906088492&u=%2Fscholar%3Fq%3Dinfo%3Aa71uJS9PPhkJ%3Ascholar.google.com%2F%26output%3Dcite%26scirp%3D0%26hl%3Den
- Sinka, M. E., Bangs, M. J., Manguin, S., Rubio-Palis, Y., Chareonviriyaphap, T., Coetzee, M., Mbogo, C. M., Hemingway, J., Patil, A. P., Temperley, W. H., Gething, P. W., Kabaria, C. W., Burkot, T. R., Harbach, R. E., & Hay, S. I. (2012). A global map of dominant malaria vectors. *Parasites & Vectors*, 5(1), 69.
<https://doi.org/10.1186/1756-3305-5-69>
- WHO. (2020). *Pictorial identification key of important disease vectors in the WHO South-East Asia Region*. World Health Organization. Regional Office for South-East Asia. <https://apps.who.int/iris/handle/10665/332202>
- World Health Organization. Regional Office for South-East Asia. (2014). *Vector-borne diseases (SEA-CD-300)*. WHO Regional Office for South-East Asia.
<https://apps.who.int/iris/handle/10665/206531>

PHOTOPLATES





Latitude: 28.674546
Longitude: 80.638997
Elevation: 196.49±7 m
Accuracy: 9.7 m
Time: 02-08-2024 07:16
Note: jugeda C2

Powered by NoteCam



Latitude: 28.676365
Longitude: 80.639378
Elevation: 202.86±31 m
Accuracy: 26.5 m
Time: 02-08-2024 18:44
Note: jugeda light trap

Powered by NoteCam

Annex: 1

Abdominal conditions of indoor resting mosquitoes from animal and human dwelling collection

S.N.	Species	Male	Female	Total	Unfed	Fulfed	Gravid	Half-gravid
Animal Dwelling								
1	<i>Anopheles fluviatilis</i>	0	9	9	0	4	3	2
2	<i>Anopheles annularis</i>	0	5	5	0	5	0	0
3	<i>Anopheles culicifacies</i>	0	1	1	0	0	0	1
4	<i>Culex quinquefasciatus</i>	9	31	40	5	16	5	5
5	<i>Armigeres subalbatus</i>	0	1	1	0	0	1	0
Human Dwelling								
1	<i>Anopheles fluviatilis</i>	0	2	2	0	1	1	0
2	<i>Anopheles annularis</i>	0	1	3	0	1	0	0
3	<i>Culex quinquefasciatus</i>	0	18	18	5	7	4	2

Annex: 2

Abdominal conditions of indoor resting mosquitoes from light trap collection

S.N.	Species	Male	Female	Total	Unfed	Fulfed	Gravid	Half-gravid
1	<i>Anopheles fluviatilis</i>		2	2		1		1
2	<i>Anopheles annularis</i>		1	1	1			
3	<i>Culex quinquefasciatus</i>	9	15	24	12	1	2	
4	<i>Culex tritaeniorhynchus</i>		2	2		2		
5	<i>Armigeres subalbatus</i>		1	1				1

Annex: 3

GPS points of houses (IRC)

IRC points	longitude	latitude
CD1	80.63961	28.67592
CD2	80.63965	28.67611
CD3	80.64068	28.67616
CD4	80.64071	28.67463
HD1	80.63953	28.67632
HD2	80.64133	28.67591
HD3	80.64106	28.67609
HD4	80.63814	28.67653
HD5	80.64114	28.67579
HD6	80.63813	28.67633

(CD=Cattle Dwelling, HD=Human Dwelling)